

Subject Index

- A**coustic neuroma. *See also*
 Neurofibromatosis type II
 hyperpigmentation and, 153, 161, 164
- Acquired deafness
 in perinatal period, 199–200
 in postnatal period, 200–201
 in prenatal period, 197, 199
- Albinism. *See also* X-linked ocular albinism
 gene for, 2
 and hearing loss, 11
- Albinism-deafness syndrome (ALDS)
 locus for, 22
 penetrance in, 20
 X-chromosome locus of, 33, 43
- ALDS. *See* Albinism-deafness syndrome
- Alport syndrome (ATS), 20, 184
 collagen IV defect and, 45
 gene for, 2, 45
 genetic mapping of, 180–182
 locus for, 22
 vestibular dysfunction and, 76
 X-chromosome locus of, 33, 43
- ALPP. *See* Placental alkaline phosphatase locus
- American Sign Language (ASL), 210, 212, 232
- American Society for Deaf Children, 215
- American Society of Human Genetics, 223
- Americans with Disabilities Act (ADA), 234
- Annual Survey of Hearing Impaired Children and Youths*, 206–207, 293–294
- Apert syndrome, digit malformation and, 75
- Ataxia, in USH1, 169–173
- ATS. *See* Alport syndrome
- Auditory nerve damage, in NF-2, 36
- Auditory system diffuse damage, in NPC, 314
- Autosomal dominant deafness inheritance, 310–312
 evidence for, 20–21
 in MFD1, 34
 in NF2, 305
 in OAV, 34
 penetrance in, 21
 in Stickler syndrome, 217–218
 in WS, 157
- Autosomal recessive deafness inheritance, 23, 25–26
 evidence for, 21–26
 incidence of, 93
 in NPC, 313
 in PDS, 23
 in USH, 167
- Autosomal recessive mutations, and rodent obesity, 102–108
- B**ell, A. G., and eugenics movement, 7–8, 10, 237
- Bilateral acoustic neurofibromatosis. *See* Neurofibromatosis type II
- Branchiootic syndrome locus, on chromosome 8, 33
- Branchiootorenal syndrome (BOR) linkages, on chromosomes 1 and 8, 288–289
- C**A-repeat polymorphisms loci, 39–40
- Carpenter syndrome, 133
- Center for Assessment and Demographic Studies (Gallaudet University), 293
- Central neurofibromatosis. *See* Neurofibromatosis type II
- Centre d'Etude du Polymorphisme Humain (CEPH), 39
 chromosome 10 linkage map from, 41
 data base families from, 285
 reference families, 40
- CH. *See* Congenital hypothyroidism
- Charcot-Marie-Tooth disease (CMT)
 autosomal dominant form of, 40
 chromosome X locus of, 33
 gene chromosome 17 locus of, 40
 genetic heterogeneity in, 40
 linkage analysis of, 41–42
 mental retardation-deafness linkage with, 182–183
- Childhood deafness in Southern Africa
 acquired, 290
 genetic syndromes in, 291
- Chondrodysplasia, in embryogenesis of cochlea, 259–261
- Chondrodysplasia punctata (CPXR), 183
 chromosome X locus of, 33
- Chondrogenesis initiation, TGF- β 1 role in, 256–258
- Chromosomal location, of human hearing-loss-linked genes, 57–58
- CMT. *See* Charcot-Marie-Tooth disease
- Cochlear ultrastructural degeneration
 in Bronx waltzer mouse, 270–273
 in chondrodysplasia, 259–261
- Cochleosaccular pathology, 69
 fluid homeostasis failure and, 70
 mutant mice physiological responses in, 72
 Reissner's membrane collapse in, 71
- COL1A. *See* osteogenesis imperfecta
- Communication
 and deaf community, 233–234
 and genetic counseling, 210, 221, 223, 227–229
- Competence, in inner ear development, 116–117

Computed tomography (CT), in deformities assessment, 69-70
 Conductive type deafness (DFN3) locus, 43
 Congenital hypothyroidism (CH), vestibular damage in, 274-276
 Congenital sensorineural deafness, sex-linked, 19
 Craniosynostosis, syndrome delineation for, 133, 135
 CT. *See* Computed tomography
 Cystic fibrosis gene, positional cloning and, 102

Deaf children, surveys of, 16-39
 Deaf community
 assessment of families from, 308-309
 and communication, 232-234
 and Gallaudet University, 233
 members of, 233
 nondirective genetic counseling for, 221
 social arrangements and, 231-232
 Deaf marriages
 affected offspring percentages from, 9
 separation/divorce prevalence in, 10
Deaf Mutism, 8-9
 Deafness. *See also* Hearing impairment;
 Hearing loss
 as a culture, 236-239
 pigmentation defects and, 152-164
 Deafness-associated defects. *See also* Specific
 defects
 range of, 72-77
 Deafness Research Foundation, 2
 Decade of the Brain, 1
 Deletion/complementation map, 88-89
 DFN3. *See* Conductive type deafness
 Diabetes-obesity syndromes, in rodents, 102
 Digit malformations
 and deafness, 73, 75
 syndromes associated with, 75
 Disomy, uniparental, 226-227
 Duchenne's muscular dystrophy, positional
 cloning and, 102
 Dystopia canthorum, in WS1, 143, 146, 149,
 155-156, 158

Endocochlear potential (EP), in
 cochleosaccular abnormalities,
 70-71
 Endolymphatic sac development, neural
 induction and, 121-122
 Epibulbar dermoids, 34
 Ethical issues, in genetic study of deafness,
 236-239
N-ethyl-*N*-nitrosourea (ENU)
 as germline point mutagen, 4
 lethal mutation induced by, 83-85, 87

Etiologic heterogeneity, 134-136
 Eugenics movement, 7, 10-11
Everyone Here Spoke Sign Language, 231
 Eye disorders, and deafness, 73

Fabry disease, 183-184
 Family assessment for linkage studies, of WS
 and US, 293-294
 Fibroblast growth factor (FGF), in
 chondrogenesis initiation, 256-258
 Fragile-X syndrome, 19

Gallaudet University, 212, 218
 Genetic Services Center, 212-217, 220, 317
 Gene-environment interaction, hearing loss
 caused by, 225-226

Gene map
 for human chromosome 10, 40
 for protein markers, 40

Gene mapping
 CA-repeat polymorphisms in, 39-40
 and genetics of hearing, 49-66
 and inherited deafness syndromes, 38-45
 LINKAGE program in, 39-40
 protein markers in, 40
 RFLPs in, 40
 VTNRs in, 40

Genetic anatomy of hearing, 32-37
 auditory nerve, 36-37
 external ear, 33-34
 inner ear, 35-36
 middle ear, 34-35
 Genetic counseling
 approaches to, 223-229
 and ASL, 210, 212, 232
 benefits of, 215
 in childhood deafness, 194
 clinical genetics in, 223-225
 communication in, 227-229
 and communication individualization, 210
 as communication process, 221, 223
 complex pedigrees analysis and, 217-220
 in congenital rubella syndrome, 217
 cultural-linguistic influences on, 213-214
 of the deaf, 212-221, 317-318
 for deaf children and adults, 214-215
 deaf couples and, 237
 deafness etiology importance in, 201, 224
 general considerations in, 216-217
 HLA typing in, 303-304
 in Huntington's disease, 228-229
 model program for, 317-318
 nondirective, 221
 recurrence risk and, 203-210, 224
 syndrome delineation significance in, 141
 for USH, 43
 USH2 linkage in, 172-173

- Genetic deafness
 autosomal dominant inherited, 193
 autosomal recessive inherited, 193-194
 heterozygous expression of, 26
 incidence of, 191-195
 X-linked, 191, 193
- Genetic epidemiology, animal models for, 27-28
- Genetic hearing impairment, frequency of, 16-18
- Genetic mapping
 in chimpanzees, 277-280
 of loci affecting hearing, 82-90
 of mouse chromosome 7 microclones, 94-95
 strategies for unmapped disorders, 184-187
 of USH2 locus, 284-287
 of WSI, 59, 66
- Genetic nosology, 133-134
- Genetic Services Center. *See* Gallaudet University
- Genetics of hearing, and genome duplication, 56-66
- Genetics of hearing impairment, history of, 6-13
- Genome duplication, and genetics of hearing, 56-66
- Genomic imprinting, and hearing impairment, 226
- Germ line mosaicism, parental (somatic), 227
- Goldenhar syndrome (GHS). *See* oculoauriculovertebral sequence
- Goodman type acrocephalopolysyndactyly. *See* Carpenter syndrome
- GT repeats, in identifying polymorphisms, 106
- Guinea pig, as genetic hearing impairment model, 11
- H**earing impairment. *See also* Deafness; Hearing loss
 causes of acquired, 16-17
 chronic middle infections and, 226
 frequency of, 16-20
 gene list of disorders with, 33
 genetic epidemiology of, 16-39
 heterogeneity of, 133-142
 late onset of, 27-28
 in 1p+ monosomy, 283
 pigmentation defects and, 164
 in pigmented rats, 274-276
 in 6q partial monosomy, 283
 in trisomy 18, 281-282
 in Turner's syndrome, 283
 as USH1-USH2 discriminator, 170, 172
- Hearing impairment genetics, history of, 6-13
- Hearing impairment in cmd/cmd mice and cartilage matrix deficiency (CMD), 265-267
 Organ of Corti defects in, 266-267
 ultrastructure studies of, 265-267
- Hearing impairment in cpk mutant mice
 molecular-structural basis of, 262-264
- Hearing impairment inheritance
 autosomal dominant, 18
 autosomal recessive, 18
 MR in, 19
 sex linkage in, 18-19
- Hearing loss. *See also* Deafness; Hearing impairment
 etiology, 315-316
 sociological implications of, 230-235
- Hearing loss studies
 paradigms for, 230
 sociological model for, 231-235
- Hemifacial microsomia. *See* oculoauriculovertebral sequence
- Hereditary deafness, mouse-human similarities of, 68-77
- Hereditary Deafness Newsletter of America*, 292
- Heterogeneity, in genetic nosology, 133-134
- Heterozygous expression, in deafness, 26
- Histidinemia mouse, as hearing impairment model, 12
- History, of genetics of hearing impairment, 6-13
- Holoprosencephaly
 etiologic heterogeneity of, 135
 syndrome delineation for, 133, 135-136
- Homeobox genes, in neural developmental regulation, 129-132
- Homeodomains, 129, 131
- Homologies, in human-mouse loci, 69, 77
- Human leucocyte antigen (HLA) typing
 in genetic counseling, 303-304
 for sensorineural hearing impairment, 303-304
- Human-on-mouse comparative gene map, 50-56
- Hunter syndrome (IDS), 20
 genetic mapping of, 184
 locus for, 22
- Huntington's disease, genetic counseling in, 228-229
- Hyperbilirubinemia, as cause of childhood deafness, 199-200, 290
- Hyperuricemia, 183
- Hypoxia, as cause of childhood deafness, 199

IDS. *See* Hunter syndrome

Induction system, in inner ear development, 116-118, 121-123, 125-127

Inductive tissue interaction, growth factor mediation of, 127

Infection in late onset hearing loss, 27

Inner ear defects, 35-36
categories, 69

mouse-human similarities in, 69

Inner ear development, factors controlling, 116-127

Inner ear malformation, in MOS transgenic mice, 268-269

Inner ear structure, mouse-human similarities in, 68-69

Inquiries into the Human Faculty, 7

Insertional mutagenesis, 130

in transgenic mouse, 131-132

Kabuki makeup syndrome, 138-139

Kearns Sayre syndrome (KSS),
mitochondrial inheritance of, 301

Kernicterus, as cause of childhood deafness, 199-200

Known-genesis syndromes, in syndrome delineation, 137, 140-141

KSS. *See* Kearns Sayre syndrome

Labyrinthine abnormalities, mouse mutant models for, 12

Labyrinthine structure assessment, by CT, 69

Language, and deaf community, 232-234

Linkage map, of human chromosome 10, 41

LINKAGE program, for gene mapping, 39

Loci affecting hearing, in humans, 54

Loci affecting mouse hearing, and human homologue location, 57-58

Locus heterogeneity, and genetic mapping, 40

Macrostomia, 34

Magnetic resonance imaging (MRI), in
Usher syndrome, 169, 172

Malar hypoplasia, 35

MANB. *See* mannosidosis

Mandibular hypoplasia, 34-35

Marriages of the Deaf in America, 9-10

MELAS. *See* Mitochondrial encephalopathy + lactic acidosis + strokelike episodes

Memoir upon the Formation of a Deaf Variety of the Human Race, 7-8

Meningiomas, familial, gene locus for, 37

Meningitis, as cause of childhood deafness, 199-201, 290

Mental retardation (MR), in inherited hearing impairment, 19

MERRF. *See* Myoclonic epilepsy with ragged red fibers

MFD1. *See* Treacher-Collins syndrome

Microphthalmia, 34

Microtia, 34-35

Middle ear, defects of, 34-35

Mitochondrial encephalopathy + lactic acidosis + strokelike episodes (MELAS), mitochondrial inheritance of, 301

Mitochondrial inheritance
of acquired deafness, 301-302

of hearing impairment, 225

of KSS, 301

of MERRF, 301

Molecular genetics, of X-linked hearing impairment, 176-187

Morbid Anatomy of the Human Genome map, 32

Morphogenetic defects, 69-70

asymmetry in, 70

CT in classification of, 70

Michel/Mondini type, 70

Morphological specific-locus test, 82, 87

Mouse autosomal recessive deafness loci,
and human homologous genes, 98

Mouse models

of genetic hearing impairment, 11-13

of hereditary deafness, 80-90

relevance to human disease of, 11-13, 101

Mouse obesity syndromes, 101-108

MR. *See* mental retardation

MRI. *See* Magnetic resonance imaging

Mutant mouse lines, genetic loci in, 80-90

Mutation identification, gene maps in, 58-59

Myoclonic epilepsy with ragged red fibers (MERRF), mitochondrial inheritance of, 301

National Association of the Deaf, 215

National Information Clearinghouse, of
NIDCD, 1

National Institute on Deafness and Other
Communication Disorders
(NIDCD), 1

National Institutes of Health, 2

National Research Register for Hereditary
Hearing Loss, 292

National Research Registry for Hereditary
Hearing Loss, 2

National Symposium on Genetic Services for
Underserved Populations, 213

NDP. *See* Norrie disease

Nerve deafness, and CMT, 40

Neural factor, in inner ear development, 118,
120

Neural tube anomalies, and deafness, 73

Neuroepithelial pathology, 69

- Neuroepithelial pathology, vestibular dysfunction in, 72
- Neurofibromatosis (NF), positional cloning and, 102
- Neurofibromatosis type II (NF2)
auditory nerve damage in, 36-37
chromosome 22 locus of, 33
gene locus for, 2, 37
treatable deafness cause, 305-307
- Neurofibromatosis type I (NF1), 305
chromosome 17 locus of, 33
gene for, 2
- Neurosensory relationships, in Bronx waltzer mouse cochlea, 270-273
- NF. *See* Neurofibromatosis
- NIDCD. *See* National Institute on Deafness and Other Communication Disorders
- Niemann-Pick type C (NPC) disease, audiologic profile in, 313-314
- Noise pollution, in late onset hearing loss, 27, 29
- Non-Mendelian inheritance, 225-227
in human obesity, 108
- Nonsyndromal genetic deafness, 191-195
- Norrie disease (NDP), 20, 33, 177-178, 180
chromosome X locus of, 33
genetic mapping of, 178, 180
locus for, 22
- NPC. *See* Niemann-Pick type C disease
- O**AV. *See* oculoauriculovertebral sequence
- Obesity, genetic basis for, 100-114
- Obesity genes
human pedigree segregating, 113
in humans, 112-114
- Oculoauriculovertebral sequence (OAV), 34
chromosome 7 gene locus of, 34
clinical features of, 34
- OI. *See* Osteogenesis imperfecta
- OMIM. *See* On-line Mendelian Inheritance in Man (OMIM)
- On-line Mendelian Inheritance in Man (OMIM), 32
- OPD. *See* otopalatodigital syndrome
- Organ of Corti
chondrodysplastic changes in, 260-261
in cpk mutant mice, 262-264
defects in, 36
degeneration, 268-269
loss in WS, 35
- Organomegaly, in NPC, 313
- Orofacial clefting, syndrome delineation for, 133, 135
- Osteogenesis imperfecta (OI), 33
autosomal dominant inheritance of, 240-247
chromosome 7 locus for, 33
collagen defect in, 35
diagnosis of, 240-241
diagnostic criteria for, 245
segregation-penetrance analysis in, 242-247
- Otic capsule, 35
- Otic placode formation, in inner ear development, 116-118
- Otocyst organogenesis, in inner ear development, 118, 120-122
- Otopalatodigital syndrome (OPD), 22
sex-linked deafness in, 20
- Otopathology evaluation, in transgenic mutant mouse, 249-251
- Otosclerosis
procollagenase mRNA amounts in, 319-321
TIMP amounts in, 319-321
- Ototoxic drugs
as cause of childhood deafness, 199, 201
cochlea damage by, 201
- P**aralogous genes, 56
mutations in, 59, 66
- Pathogenetic heterogeneity, 134-136
- PBT. *See* Piebaldism
- PCR. *See* polymerase chain reaction
- PDS. *See* Pendred syndrome
- Pendred syndrome (PDS), 23, 33
in childhood deafness, 291
- Penetrance, in albinism/deafness syndrome, 20
- Perinatally acquired deafness
hypoxia as cause of, 199
kernicterus, 199-200
- PFG. *See* Pulsed-field gel electrophoresis
- Piebaldism (PBT), and WS, 158-159, 164
- Pigmentation defects
and deafness, 72-74, 152-164
in WS, 73-74, 143-144
- Pinna defects, 33-34
and deafness, 73-74
- Placental alkaline phosphatase locus (ALPP), lod scores with WS of, 147-149
- Pleiotropy, in genetic nosology, 72, 133-134
- Polycystic kidney mice, inner ear ultrastructure in, 262-265
- Polygenic inheritance, 108-112
- Polymerase chain reaction (PCR), 94, 97, 131-132
in marker typing, 39
- Positional cloning
and cystic fibrosis gene, 102
and Duchenne's muscular dystrophy, 102
technique, 102-103
- Postnatally acquired deafness
meningitis as cause of, 198, 200-201
ototoxic drugs as cause of, 201, 290

- POU specific domain gene expression, in developing rat cochlea, 129-131
- Prelingual hearing impairment, prevalence of, 16
- Prenatally acquired deafness
 rubella as cause of, 197-199, 290
 syphilis as cause of, 199
- Presbycusis
 diet effect on, 13
 in late onset hearing loss, 27, 29
- Program of Services to Deaf Patients, 308
- Protein markers, on gene maps, 40
- Protooncogene int-2, as inductor candidate, 127
- Pulsed-field gel (PFG) electrophoresis, 89
- Q**uantitative trait loci (QTL), 111-112
- R**adiation-induced deletion mutations, 87, 89
- Recombinant DNA Advisory Committee, of NIH, 2
- Recurrence risk
 in Alport syndrome, 205
 factors in calculation of, 203
 implications for genetic counseling, 203-210
 for isolated deafness, 209-210
 in known genetic conditions, 203-206
 loci abundance implications for, 20
 SEGRAN analysis of, 206-208
 when cause unknown, 206
 in WS, 203-205
- Recurrent-pattern syndromes, in syndrome delineation, 137, 139-141
- Rehabilitation Act of 1973, 234
- Reissner's membrane, in cpk mutant mice, 262-264
- Reissner's membrane collapse
 in cochleosaccular pathology, 71
 in fluid homeostasis failure, 70
- Reproduction, in deaf populations, 26-27
- Restriction fragment length polymorphism (RFLP), 97, 103, 106, 111, 114, 144, 184, 292, 295
 DNA markers and, 38-39
 Southern blot test, 39
 human obese phenotype linkage to, 101
 loci, 40
- Restriction fragment length variant (RFLV), 85
 in gene mapping, 94-95
- Retinitis pigmentosa (RP)
 chromosome 3 locus of, 33
 chromosome 3 mutation in, 3
 rhodopsin gene mutation in, 3
 and US, 167-168
 as USH1-USH2 discriminator, 172
- Reverse genetics
 and deafness gene cloning, 93-98
 and deafness studies, 80-90
- RFLP. *See* Restriction fragment length polymorphism
- RFLV. *See* Restriction fragment length variant
- Rhombencephalon, in inner ear development, 118, 120-121, 123, 125-126
- Rubella (congenital), as cause of childhood deafness, 197-199, 201
- S**cheibe-type abnormality. *See* Cochleosaccular pathology
- Sclerosteosis, in childhood deafness, 291
- Segregation frequencies, for sporadic cases, 22
- SelfHelp for Hard of Hearing, Inc., 215
- Sensorineural hearing impairment, HLA frequencies in, 303-304
- Sewall Wright formula, for F2 phenotypic extremes, 110-111
- Sex-linked deafness
 albinism-deafness syndrome, 20
 conductive type with gusher, 19
 congenital sensorineural type, 19
 Hunter syndrome (IDS), 20, 22, 184
 mutation rate estimates, 20
 Norrie disease (NDP), 22, 178, 180
 otopalatodigital, syndrome (OPD), 22
 pedigrees of, 19-20
 Toriello syndrome, 184-187
- Shaker-1, as genetic hearing impairment model, 12
- Sign language
 cultural basis of, 236-237
 and deaf community, 232-234
- Single gene mutation, in external ear formation, 33
- Skeletal abnormalities, and deafness, 73
- Social arrangements, and deaf community, 231-232
- Spastic paraplegia, genetic heterogeneity in, 40
- Stapes fixation with gusher (DFN3), locus for, 22
- Stickler syndrome, pedigree analysis of, 217-218
- Stria vascularis dysfunction, melanocyte absence and, 72
- Summitt syndrome. *See* Carpenter syndrome
- Syndrome delineation, 133-142
 involving craniostylosis, 133, 135
 involving holoprosencephaly, 133, 135-136
 involving orofacial clefting, 133, 135
 significance of, 141-142
 stages of, 136-137, 139-140

Synotia dorsalis, experimental production of, 121

Syphilis (congenital), as cause of childhood deafness, 199, 290

Telecommunication device for the deaf (TDD), 218, 231, 234

Thalidomide, and deafness, 201, 290

Toriello syndrome, 184-187

Transforming growth factor-beta ($TGF-\beta_1$) chondrogenesis initiation role of, 256-258 in inner ear morphogenesis, 253-255

Treacher-Collins syndrome (MFD1) in childhood deafness, 291 chromosome 5 gene locus of, 34 clinical features of, 35

Tuberous sclerosis, genetic heterogeneity in, 40

Unknown-genesis syndromes, in syndrome delineation, 136-137, 139, 141

USH. *See* Usher syndrome

Usher syndrome (USH), 23

ataxia in, 169-173

autosomal recessive inheritance of, 167

in childhood deafness, 291

genetic heterogeneity within, 43

genetics of, 163-164

genetic studies of, 167-173

history and prevalence of, 162

identity by descent in, 26

retinal pigmentation and, 161-162

type distinction of, 167-169

Usher syndrome type I (USH1)

clinical characteristics of, 43

linkage analysis vs. chromosome 3

markers, 298-300

location score for, 44

Usher syndrome type II (USH2)

chromosome 1 locus of, 33, 42

clinical characteristics of, 43

gene for, 2

in genetic counseling, 172-173

localization to chromosome 1q of, 284-287

location of, 170-173

locus for, 22

marker THH33 linkage with, 43-44

RP in, 284

vestibular system involvement in, 284

Van der Hoeve's triad, collagen defect in, 35

Variability in number of tandem repeats (VTNR)

loci, 40

markers, 39

Vestibular damage, in CH, 274-276

Vestibular dysfunction

and deafness, 73, 75-77

in neuroepithelial pathology, 72

syndromes associated with, 76

Vestibular function, as USH1-USH2 discriminator, 170

Vitamin D-resistant rickets. *See* X-linked hypophosphatemia

Von Recklinghausen's disease. *See* Neurofibromatosis type I (NF1)

VTNR. *See* Variability in number of tandem repeats (VTNR)

Waardenburg syndrome (WS), 35-36

in childhood deafness, 291

diagnostic criteria for, 157

families assessment for, 308-309

gene locus for, 36

genetic mapping of, 45

genetics of, 157-158

history and nosology of, 153-157, 164

mutation rate for, 21

white spotting in, 71

Waardenburg syndrome type I (WS1)

ALPP locus linkage with, 147-148

chromosome 2 locus of, 33

clinical manifestations of, 143-144

exclusion mapping of, 146-147

localization of gene for, 143-149

localization on chromosome 2q, 295-297

locus for, 22, 55

spotch homology with, 55

White spotting, in deafness syndromes, 71, 73, 143-144

WS. *See* Waardenburg syndrome

X-linked albinism-deafness syndrome (ADFN), blood clotting factor 9 linkage with, 183

X-linked deafness

forms of, 176-178

molecular genetics of, 176-187

X-linked hypophosphatemia (Hyp), genetic mapping of, 184

X-linked ocular albinism with deafness (XLASD), 159-161

X-linked progressive mixed deafness + perilymphatic gusher (DFN3), 178

X-linked spastic paraplegia, 40

Yeast artificial chromosomes (YAC), 89, 97-98, 103-104, 108